

TECHNICAL NOTE

Measurements of cyclic AMP and cyclic GMP phosphodiesterase activity in isolated tubular segments

BRIAN A. JACKSON, RICHARD M. EDWARDS, and THOMAS P. DOUSA
with the technical assistance of DENISE M. HEUBLEIN and JULIE L. BRAUN

Nephrology Research Laboratory, Department of Physiology & Biophysics and Division of Nephrology, Mayo Clinic and Foundation, Rochester, Minnesota

The structural and functional heterogeneity of the renal tubular system is well recognized. To date however, the biochemical steps in the mechanism of action of hormones whose effects are mediated by cyclic 3',5'-AMP (cAMP) have been studied primarily in the whole kidney or in major anatomic subdivisions (cortex, medulla, and papilla) of mammalian renal tissue [1, 2].

Recently, however, Imbert, Chabardes, and Morel introduced a micromethod for the assay of adenylate cyclase activity in specific nephron segments [3, 4] and demonstrated that adenylate cyclase stimulated by vasopressin (VP), parathyroid hormone, calcitonin, or isoproterenol is limited to well-defined segments of the renal tubular system [5].

Stimulation of adenylate cyclase, although important, is only the initial step in the sequence of intracellular biochemical events leading to the ultimate functional response of the tubule to a hormone. The rate of cAMP accumulation and the intracellular cAMP concentration are dependent not only on the rate of biosynthesis but also on the rate of hydrolysis by a specific enzyme, cyclic 3',5'-AMP phosphodiesterase (cAMP-PDIE), and on the rate of efflux of cAMP from the cell [6]. Our previous studies [7] indicated that the distribution of enzymes of cAMP and cyclic 3',5'-GMP (cGMP) metabolism were different in glomeruli compared to mixed cortical tubular fractions or to cortical slices, suggesting that the catabolism of cyclic nucleotides differs in specific substructures of the nephron.

Cyclic nucleotide catabolism to date has not been studied in specific segments of the renal tubular system, except for a semiquantitative histochemical study of cAMP-PDIE in rabbit tissues, including kidney, which suggested the activity of this enzyme differed in various tubular segments [8]. In particular, high cAMP-PDIE activity was reported in the ascending limb of the loop of Henle, but little or no activity was reported in the collecting tubule. We have therefore developed a method for the determination of both cAMP-PDIE and cGMP phosphodiesterase (cGMP-PDIE) activity in specific tubular segments microdissected from the rat kidney. In particular, we have applied this technique to measure the activity of these two enzymes in the medullary collecting tubule (MCT) and the medullary thick ascending limb of the loop of Henle (MAL), two nephron segments recently shown to be major sites of VP-sensitive adenylate cyclase activity [3, 5, 9] and responding to VP and cAMP by changes in water permeability [10-13] or solute transport [14].

Methods and results

Adult male Sprague-Dawley rats (body wt, 200 to 250 g) were allowed free access to both food and water prior to sacrifice. Outer medullary tissue was prepared for microdissection by a method described by Imbert and Morel et al [3, 4] with several modifications. Briefly, while the rats were under light pentobarbital anesthesia (6 mg/100 g of body wt), the aorta directly above the left renal artery was clamped and the left kidney perfused with 15 ml of heparinized (20 USP units/ml) "collagenase medium" (for composition, see below) at a rate of 1.5 ml/min. Corticomedullary tissue slices were incubated in this medium for 30 min at 35° C, then washed thoroughly by rinsing in "microdissection medium"

Received for publication April 19, 1979
and in revised form December 5, 1979

0085-2538/80/0018-0512 \$01.40

© 1980 by the International Society of Nephrology

Table 1. Adenylate cyclase activity sensitive to [8-Arginine]-vasopressin (AVP) and salmon calcitonin (SCT), in rat medullary thick ascending limb of Henle's loop (MAL) and in medullary collecting tubule (MCT)^a

	MAL	<i>P</i>	MCT	<i>P</i>
Basal	12.4 ± 1.2 (5)		31.7 ± 5.2 (5)	
AVP (10 ⁻⁶ M)	322.3 ± 23.0 (4)	<0.001	366.0 ± 49.0 (5)	<0.001
SCT (0.1 µg/ml)	160.2 ± 23.9 (5)	<0.001	30.0 ± 4.8 (4)	NS

^a Values for specific activity of adenylate cyclase (mean ± SEM) are expressed in fmoles/30 min/mm of tubule length. Numbers in parentheses denote the number of samples. *P* values are for significance of difference between basal and hormone-stimulated activities (group *t* test).

(137 mM sodium chloride, 5 mM potassium chloride, 0.8 mM magnesium sulfate, 0.33 mM dibasic sodium phosphate, 0.44 mM monobasic potassium phosphate, 1.0 mM magnesium chloride, 10 mM Tris hydrochloric acid, 0.25 mM calcium chloride; pH, 7.4) and transferred to Petri dishes for microdissection. Subsequent preparative procedures were performed at 0 to 4° C. (Composition of "collagenase medium" was identical to "microdissection medium" except that calcium chloride concentration was 1 mM (instead of 0.25 mM); and 0.075% collagenase (Type I, 125-250 U/mg; Worthington Biochem. Corp., Freehold, New Jersey), 0.1% hyaluronidase (Type I, Sigma Chem. Co., St. Louis, Missouri), and 0.1% bovine serum albumin was added.

Individual tubular segments (0.5 to 2.0 mm in length) were dissected under stereoscopic magnification with sharpened stainless steel needles and transferred in a small volume (less than 1 µl) of "microdissection medium" by micropipette onto concave bacteriologic slides. All segments were then photographed for subsequent length measurement [4, 9]. The identity of each segment established by visual inspection was confirmed from the photograph by distinct morphologic characteristics [3, 4]. In addition, adenylate cyclase responses to VP (10⁻⁶ M [8-Arg]-vasopressin) and to synthetic salmon calcitonin (SCT; 0.1 µg/ml) were tested in a preliminary experiment (Table 1). Segments identified as ascending limb of Henle's loop (MAL) contained adenylate cyclase sensitive to both VP and SCT, whereas segments of medullary collecting tubules (MCT) contained only VP-sensitive adenylate cyclase, an observation consistent with that of Morel, Chabardes, and Imbert-Teboul [5] on other species.

Tubular samples were "permeabilized" by aspirating off the excess microdissection medium and

Table 2. Effect of freezing and thawing on cAMP-phosphodiesterase (cAMP-PDIE) activity in medullary thick ascending limb of Henle's loop (MAL) and medullary collecting tubule (MCT)^a

	MAL	<i>P</i>	MCT	<i>P</i>
Nonfrozen	8.7 ± 0.8 (5)		5.0 ± 1.0 (4)	
Frozen	16.9 ± 2.6 (5)	<0.02	19.8 ± 1.1 (5)	<0.001

^a Values for cAMP-PDIE activity (mean ± SEM) are expressed in fmoles/min/mm of tubule length. Numbers in parentheses denote the number of samples. *P* values are for significance of difference between frozen and nonfrozen activities (group *t* test).

replacing with 0.5 µl of "hypotonic medium" (1 mM magnesium chloride, 0.25 mM ethylene-diamine-tetracetic acid (EDTA), 0.1% bovine serum albumin (wt/vol), and 1 mM Tris hydrochloric acid; pH, 7.4), followed by rapid freezing on dry ice. Omission of this freezing and thawing step results in a significant reduction in cAMP-PDIE activity in both MAL and MCT, as observed in a preliminary experiment (Table 2). A similar observation was made with respect to adenylate cyclase activity measurements by Morel et al [4] and was attributed to an inability of enzyme substrate to enter the cell unless permeabilized by freezing and thawing. All samples were stored overnight at -80° C prior to assay. No significant loss of enzyme activity occurred in either MCT or MAL stored overnight at -80° C when compared with samples assayed the same day (Table 3).

Cyclic AMP-PDIE and cyclic GMP-PDIE activities were assayed by incubating permeabilized tubular segments in 5 µl of a medium of the following composition: 10 mM magnesium sulfate, 0.1 mM EDTA; 50 mM Tris hydrochloric acid; pH, 8.0 (Mg-EDTA-Tris buffer). Final substrate concentration (either ³H-cAMP or ³H-cGMP; New England Nuclear Co., Boston, Massachusetts) was 10⁻⁶ M (3 × 10⁵ cpm per reaction mixture). After addition of the incubation mixture to the tubule, the slide was covered with a second vaseline-coated slide, providing a waterproof reaction chamber, which was immersed in a water bath maintained at 37° C for the desired length of time. At the end of the incubation period, 50 µl of Mg-EDTA-Tris buffer was added to the reaction mixture, and the mixture was transferred by micropipette into glass test-tubes (12 × 75 mm); the slide was rinsed with another 50 µl of Mg-EDTA-Tris buffer, which was added to the test tube. The test tubes were immersed for 3 min in boiling water to terminate the enzyme reaction. Fifty microliters of 5'-nucleotidase (snake venom from

Table 3. Effect of 24-hr storage at -80°C on cAMP-phosphodiesterase (cAMP-PDIE) activity in medullary thick ascending limb of Henle's loop (MAL) and medullary collecting tubule (MCT)^a

	MAL	<i>P</i>	MCT	<i>P</i>
Same day	13.0 \pm 1.0 (12)		14.3 \pm 0.9 (12)	
24-hr storage	14.3 \pm 1.0 (12)	NS	14.3 \pm 1.1 (12)	NS

^a Values for cAMP-PDIE activity (mean \pm SEM) are expressed in fmoles/min/mm of tubule length. Numbers in parentheses denote the number of samples. *P* values are for significance of difference between same day and 24-hr storage activities (group *t* test).

Crotalus atrox; 1 mg/ml) were added to each tube and incubated for a further 15 min at 37°C as in our previous studies [7, 15]. Nucleotides were separated from nucleosides on QAE-Sephadex columns [7, 15] with 3 ml of 20 mM ammonium formate (pH, 7.4). Approximately 95% recovery from QAE-Sephadex columns was achieved under present conditions, results similar to those reported by Wells et al [16]. A standard assay blank of 5 μl of the reaction mixture without tubular protein was included in every assay. For cAMP-PDIE assays, the blank value never exceeded 2% (6000 cpm) of the total cpm added, and for cGMP-PDIE, the blank value never exceeded 4% (12,000 cpm) of the total counts per minute added.

Under the present conditions, cAMP hydrolysis in both MAL and MCT was linearly proportional to time for at least 40 min (hydrolysis of about 15% of substrate) (Fig. 1), whereas hydrolysis of cGMP progressed linearly with time only up to 15 min (hydrolysis of about 10% substrate), in both segments (Fig. 1). Therefore, in subsequent experiments, an incubation time of 20 min was used for the cAMP-

PDIE assay and 10 min for the cGMP-PDIE assay. Both cAMP-PDIE and cGMP-PDIE activities were directly proportional to tubular length both in MAL and MCT over the ranges 1.0 to 7.0 mm and 1.0 to 5.0 mm, respectively (Fig. 2). Samples consisting of a total of approximately 1 to 2 mm of tubule were assayed in all subsequent experiments. Assays were conducted under conditions when activities were at least 2 to 3 times higher than blank values.

The specific activities of cAMP-PDIE and cGMP-PDIE in MAL and MCT were compared in such a way that segments of MAL and MCT microdissected from the same animal were always assayed for cAMP-PDIE or cGMP-PDIE simultaneously in order to allow paired comparison (Table 4). The specific activity of cGMP-PDIE was significantly higher ($P < 0.001$; Student's *t* test) than was cAMP-PDIE activity in both MAL and MCT. The rate of both cAMP and cGMP hydrolysis was similar in MCT compared with MAL, expressed per millimeter of tubule length (Table 4a). To compare the specific activities of cAMP- and cGMP-PDIE in isolated nephron segments with activities measured on renal homogenates and tissue extracts, we also calculated the data relative to tubular protein (Table 4b). Because it was not feasible to measure both enzyme activity and protein content on the same sample, such values could only be calculated indirectly from the actual tubular length, and a mean value for protein content for each segment, determined in separate experiments. Protein content was determined by a micromodification [17] of the micro method of Lowry et al [18]; both samples and standards contained 1% sodium dodecyl sulfate (SDS). (SDS does not interfere with the protein determination [19].) The mean protein content of MCT ($0.11 \pm [\text{SEM}] 0.01 \mu\text{g}/\text{min}$; $N = 24$) was simi-




Fig. 1. Dependence of cAMP (left panel) and cGMP (right panel) hydrolysis on the time of incubation in medullary collecting tubule (MCT; \bullet - \bullet) and in medullary ascending limb (MAL; \circ - \circ). Each point represents mean \pm SEM of four to six samples. For details see text.




Fig. 2. Dependence of cAMP-phosphodiesterase (upper panels) and cGMP-phosphodiesterase (lower panels) activity on tubular length in the incubation mixture. Left hand panels represent medullary collecting tubule (MCT), and right hand panels represent medullary ascending limb (MAL). Regression lines were fitted by the method of least squares. For details see text.

lar to that of MAL ($0.13 \pm [\text{SEM}] 0.01 \mu\text{g}/\text{mm}$; $N = 35$). When expressed per unit of protein, only minor differences in specific activities of either cAMP-PDIE or cGMP-PDIE existed between MCT and MAL (Table 4b).

Both cAMP-PDIE and cGMP-PDIE activities (Fig. 3) were almost completely inhibited by the addition of $5 \times 10^{-4} \text{ M}$ 1-methyl-3-isobutylxanthine (MIX)—a known phosphodiesterase inhibitor [1, 7, 13]—in both segments. A concentration of $5 \times 10^{-5} \text{ M}$ MIX caused a partial inhibition of both cAMP-PDIE and cGMP-PDIE in MAL and MCT.

Discussion

Results of the present studies demonstrate that the rate of cAMP and cGMP hydrolysis can be measured in specific microdissected segments of the renal tubule. Although there are probably several cyclic nucleotide phosphodiesterases in the kidney [20], this investigation was confined to one substrate concentration (10^{-6} M), which reflects predominantly, but not exclusively, the so called low K_m -high affinity enzyme activity [21]. cGMP-PDIE activity is higher than cAMP-PDIE is, in both MCT and MAL. Both cGMP and cAMP-PDIE activities

Table 4. Specific activities of cAMP-PDIE and cGMP-PDIE in rat medullary thick ascending limb of Henle's loop (MAL) or in medullary collecting tubule (MCT)^a

	N	MCT	MAL	P ^b	P ^c
(a) fmoles/min/mm tubule length					
cAMP-PDIE	(14)	16.9 ± 9.9^a	16.8 ± 0.7	NS	NS
cGMP-PDIE	(5)	52.6 ± 5.8	52.0 ± 6.0	NS	NS
(b) fmoles/min/ μg protein					
cAMP-PDIE	(14)	153.8 ± 8.7^a	129.7 ± 5.5	<0.05	<0.025
cGMP-PDIE	(5)	478.8 ± 53.6	400.4 ± 46.8	NS	<0.025

^a Numbers in parentheses (N) denote the number of animals; value of each animal is the mean of three to five samples. Values for both cAMP-PDIE and cGMP-PDIE activities (mean \pm SEM) are expressed in fmoles/min/mm of tubule length (a) and fmoles/min/ μg of protein (b).

^b Significance of difference between MCT and MAL (group *t* test)

^c Significance of difference between MCT and MAL (paired *t* test)




Fig. 3. Inhibitory effect of 1-methyl-3-isobutylxanthine (MIX) on cAMP-phosphodiesterase (top panel) and cGMP-phosphodiesterase (bottom panel) activities. Columns represent mean \pm SEM; numbers in parentheses at the bottom of columns denote number of samples. Asterisk (*) denotes values significantly different ($P < 0.05$ or higher levels of significance; t test) from the corresponding controls.

are similar in these two segments. This observation of similar cAMP-PDIE activities in MCT and MAL differs from a previously reported histochemical study on the rabbit kidney [8] in which higher activity was reportedly present in the MAL. In that study, however, in addition to the species difference, a high (10^{-3} M) concentration of cAMP was used, and evaluation of PDIE activity was only semiquantitative.

The method we have developed is essentially a scaled-down version of the standard macroassay for phosphodiesterase used frequently both in our and other laboratories for determinations in tissue homogenates and fractions [7, 14, 15, 19]. A number of dissimilarities do exist, however. For example, tech-

nical limitations make it impossible for these small tubular fragments to be homogenized, and therefore, segments are made permeable to enzyme substrate by a process of freezing and thawing in a hypotonic solution [4], which in effect causes the cells to "disintegrate." The tubules after this treatment are essentially equivalent to a "crude homogenate." The necessity for this procedure, originally designed for measuring adenylate cyclase activity [3, 4], and later for determination of ATPase activity in single segments of the nephron [22], is demonstrated in Table 2, in which untreated tubular samples show significantly lower PDIE activity. It cannot be totally excluded that disintegrated tubular cells from different segments of the nephron may be differentially permeable to substrate. However, this would be extremely difficult to test directly. Under these conditions, it is perhaps more important to ensure that both segments are treated identically. It is unlikely that freezing and thawing directly effects phosphodiesterase activity. Our laboratory data for cortical homogenates showed that freezing and thawing had no effect on either cAMP- or cGMP-PDIE activity when compared with fresh preparations (unpublished observations).

Specific activities have generally been expressed relative to tubular length, as was done in previous studies in which adenylate cyclase [3, 4] and ATPase [22] activities were measured in isolated nephron segments. Alternatively, expression of specific enzyme activities relative to tubular protein content (calculated from tubule length and mean protein content for a given segment) may be of value when comparing specific activities of cAMP-PDIE and cGMP-PDIE of microdissected tubules with other preparations such as homogenates, and possibly in comparing activities of two segments with markedly different protein content. But because the mean protein content of MCT did not differ markedly from that of MAL, specific activities expressed either per tubule length or per tubule protein were similar for these two segments (Table 4, a and b). In addition, we find that the specific activities of both cAMP-PDIE and cGMP-PDIE in MCT and MAL compare favorably with the activities of these two enzymes measured in a mixed tubular suspension from rat renal cortex [7].

In contrast to the cortex, the specific activity of cGMP-PDIE is higher than is cAMP-PDIE activity, which may be compatible with reports that cGMP metabolism [23, 24], including breakdown [23], is more active in medulla and papilla than it is in the cortical zone of the rat kidney.

Our data would suggest then, that in the normal rat kidney, cAMP formation in response to VP does not appear to be either offset or favored by a differential rate of cAMP catabolism in the MCT compared with the MAL. VP-sensitive cAMP synthesis has been reported to be impaired in renal tubular disorders, such as the urinary concentrating defect seen in Brattleboro rats with diabetes insipidus [25], or in experimental uremia in rabbits [26], or in mice with hereditary nephrogenic diabetes insipidus [27]. The importance of measuring cAMP catabolism is illustrated by our recent finding that abnormal cAMP catabolism may play a key role in the inability of MCT to increase cAMP levels in response to VP [27]. We observed that in MCT from mice with nephrogenic diabetes insipidus, cAMP-PDIE activity is abnormally high and that VP responsiveness (in terms of cAMP accumulation) could be partially restored by treatment with a phosphodiesterase inhibitor MIX [27]. Thus, use of the methodology outlined in this communication in studying other models of renal tubular disorders may similarly provide new insights into the pathogenesis of anomalous responsiveness of renal tubules to VP [25, 26, 27] and to other hormonal agents [28] whose actions are mediated by cAMP.

Acknowledgments

A preliminary report of this work was published in abstract form (*Clin Res* 26:780, 1978). This research was supported by USPHS NIH Grant AM-16105 from the National Institute of Arthritis, Metabolism and Digestive Diseases and by the Mayo Foundation. Dr. B. A. Jackson was the recipient of a Mayo Postdoctoral Research Fellowship and Wellcome Travel Award. Part of this research was conducted by R. M. Edwards in partial fulfillment of the requirements for the Ph.D. degree in Physiology, Mayo Graduate School of Medicine. Dr. T. P. Dousa, was an Established Investigator of the American Heart Association (AHA-74182). Mrs. A. M. Walker provided secretarial assistance. Dr. D. Chabardes, Dr. M. Imbert-Teboul, and Prof. F. Morel, Laboratoire de Physiologie Cellulaire, College de France, Paris, France, gave advice in establishing tubule microdissection techniques.

Reprint requests to Dr. T. P. Dousa, Mayo Clinic, Guggenheim Bldg., 921B, Rochester, Minnesota 55901, USA

References

- DOUSA TP: Drugs and other agents affecting the renal adenylate cyclase system, in *Methods in Pharmacology*, edited by
- MARTINEZ-MALDONADO M, New York, Plenum Press, 1976, vol. 4A, pp. 293-331
- DOUSA TP: Cyclic nucleotides in renal pathophysiology, in *Contemporary Issues in Nephrology*, edited by STEIN JH, BRENNER BM, New York, Churchill Livingstone Inc., 1979, vol. 4, pp. 251-285
- IMBERT M, CHABARDES D, MONTEGUT M, CLIQUE A, MOREL F: Adenylate cyclase activity along the rabbit nephron as measured in single isolated segments. *Pfluegers Arch* 354:213-228, 1975
- MOREL M, CHABARDES D, IMBERT-TEBOUL M: Methodology for enzymatic studies of isolated tubular segments: Adenylate cyclase, in *Methods in Pharmacology*, edited by MARTINEZ-MALDONADO M, New York, Plenum Press, 1978, vol. 4B, pp. 297-323
- MOREL F, CHABARDES D, IMBERT-TEBOUL M: Hormonal control of cyclic AMP generation by isolated nephron segments and its relation to problems of kidney physiology, in *Renal Function*, edited by GIEBISCH G, PURCELL E, Baltimore, Waverly Press, 1977, pp. 275-289
- SAMIR AMIR M: Cyclic nucleotides as targets for drug design, in *Advances in Drug Research*, edited by HARPER NJ, SIMMONDS AB, New York, Academic Press, 1977, pp. 1-38
- TORRES VE, HUI YSF, SHAH SV, NORTHRUP TE, DOUSA TP: Cyclic nucleotide phosphodiesterases in glomeruli of rat renal cortex. *Kidney Int* 14:444-451, 1978
- SHANTA TR, WOODS WD, WAITZMAN MB, BOURNE GH: Histochemical method for localization of cyclic 3',5'-nucleotide phosphodiesterase. *Histochemie* 7:177-190, 1966
- IMBERT-TEBOUL M, CHABARDES D, MONTEGUT M, CLIQUE A, MOREL F: Vasopressin-dependent adenylate cyclase activities in the rat kidney medulla: Evidence for two separate sites of action. *Endocrinology* 102:1254-1261, 1978
- ORLOFF J, HANDLER JS: The similarity of effects of vasopressin, adenosine 3',5'-monophosphate (cyclic AMP) and theophylline on the toad bladder. *J Clin Invest* 41:702-709, 1962
- GRANTHAM JJ, BURG MB: Effect of vasopressin and cAMP on permeability of isolated collecting tubules. *Am J Physiol* 211:255-259, 1966
- BECK NP, KANEKO T, ZOR U, FIELD JB, DAVIS BB: Effects of vasopressin and prostaglandin E₁ on the adenylate cyclase-cyclic 3',5'-adenosine monophosphate system of the renal medulla of the rat. *J Clin Invest* 50:2461-2465, 1971
- HALL DA, BARNES LD, DOUSA TP: Cyclic AMP in action of antidiuretic hormone: effects of exogenous cyclic AMP and its new analogue. *Am J Physiol* 232(4):F368-F376, 1977
- HALL DA: Possible role of vasopressin in regulating solute transport in mouse thick ascending limb of Henle's loop. *Clin Res* 27:416A, 1979
- KIM JK, FROHNERT PP, HUI YSF, BARNES LD, FARROW GM, DOUSA TP: Enzymes of cyclic 3',5'-nucleotide metabolism in human renal cortex and renal adenocarcinoma. *Kidney Int* 12:172-183, 1977
- WELLS JN, BAIRD CE, WU YJ, HARDMAN JG: Cyclic nucleotide phosphodiesterase activities of pig coronary arteries. *Biochim Biophys Acta* 384:430-442, 1975
- BRENNER BM, FALCHUK FH, KEIMOWITZ RI, BERLINER RW: The relationship between peritubular capillary protein concentration and fluid reabsorption by the renal proximal tubule. *J Clin Invest* 48:1519-1531, 1969
- LOWRY OH, ROSENBOUGH NJ, FARR AL, RANDALL RJ: Protein measurement with folin phenol reagent. *J Biol Chem* 193:265-275, 1951

19. DOUSA TP, WILSON DM: Effects of demethylchlortetracycline on cellular action of antidiuretic hormone in vitro. *Kidney Int* 5:279-284, 1974
20. VAN INWEGEN RG, SWAFFORD RL, STRADA SJ, THOMPSON WJ: Characterization of particulate cyclic nucleotide phosphodiesterase of rat kidney. *Arch Biochem Biophys* 179:58-68, 1977
21. STRADA SJ, THOMPSON WJ: Multiple forms of cyclic nucleotide phosphodiesterases: anomalies or biologic regulators? *Adv Cycl Nucl Res* 9:265-283, 1978
22. DOUCET A, KATZ AI, MOREL F: Determination of Na-K-ATPase activity in single segments of the mammalian nephron. *Am J Physiol* 237:F105-F113, 1979
23. CRAVEN PA, DERUBERTIS FR: Properties and subcellular distribution of guanylate cyclase activity in rat renal medulla: Correlation with tissue content of guanosine 3',5'-monophosphate. *Biochemistry* 15:6131-6137, 1976
24. SHAH SV, NORTHRUP TE, HUI YSF, DOUSA TP: Action of serotonin (5-hydroxy-tryptamine) on cyclic nucleotides in glomeruli of rat renal cortex. *Kidney Int* 15:463-472, 1979
25. IMBERT-TEBOUL M, CHABARDES D, MONTEGUT M, CLIQUE A, MOREL F: Impaired response to vasopressin of adenylate cyclase of the thick ascending limb of Henle's loop in Brattleboro rats with diabetes insipidus. *Renal Physiol* 1:3-10, 1978
26. FINE LG, SCHLONDORFF D, TRIZNA W, GILBERT RM, BRICKER MS: Functional profile of the isolated uremic nephron: Impaired water permeability and adenylate cyclase responsiveness of the cortical collecting tubule to vasopressin. *J Clin Invest* 61:1519-1528, 1978
27. JACKSON BA, EDWARDS RM, VALTIN H, DOUSA TP: Cellular action of vasopressin in medullary tubules of mice with hereditary nephrogenic diabetes insipidus. *J Clin Invest* 64:110-122, 1980
28. BRUNETTE MG, CHABARDÈS D, IMBERT-TEBOUL M, CLIQUE A, MONTEGUT M, MOREL F: Hormone-sensitive adenylate cyclase along the nephron of genetically hypophosphatemic mice. *Kidney Int* 15:357-369, 1979